

Astrofisica e particelle elementari

aa 2010-11

Lezione 7

- Esperimenti su satellite per radiazione gamma
- Cerenkov in atmosfera, esperimenti a terra per radiazione gamma

Bruno Borgia

Prossime conferenze

- 2011 Fermi Symposium 9-12 May 2011 Rome,
<http://fermi.gsfc.nasa.gov/science/symposium/2011/>
- 3rd Roma International Conference on Astro-Particle Physics 25-27 May Rome
<http://ricap11.roma3.infn.it/>


FERMI SYMPOSIUM

LUNEDI 9 SESSIONE PLENARIA

9:00-9:20	welcomes
9:20 - 9:45	Fermi Mission Overview
9:45 - 10:10	Fermi and Astrophysics (Steve Shore)*
10:10 - 10:35	Enrico Fermi and High Energy Astrophysics in Italy (G.F. Bignami)*
10:35 - 11:00	break
11:00 - 11:30	Fermi and Fermi (Ugo Amaldi)*
11:30 -11:55	The Second Fermi LAT Catalog: Construction and Contents (Toby Burnett)*
11:55 - 12:15	The Second Fermi LAT Catalog: Caveats and Classifications (Dave Thompson)*
12:15 - 12:30	TeV sources analysis with AGILE (Longo)
12:30 - 14:00	Lunch/Posters
14:00 - 14:30	Galactic diffuse emission: models and interpretation (Jean Marc Casandjian)*
14:30 - 14:45	Fermi's View of the Inner Galaxy (Porter)
14:45 - 15:00	Cosmic-Ray Positron Measurement with the Fermi-LAT Using the Earth's Magnetic Field (Mitthumsiri)
15:00 - 15:15	Results from the PAMELA experiment (Papini)
15:15 - 15:30	Fermi Bubbles: A 10 Kpc Shock From The Galactic Center? (Su)
15:30 - 16:00	break
16:00 - 16:30	Solar Flares and Gamma-ray Observations (Jim Ryan)*
16:30-16:45	Impulsive High-Energy Particle Acceleration in the SOL2010-06-12T00:57 M2 X-ray Flare (Gerry Share)
16:45 - 17:00	Long-lived solar gamma-ray emission during 2011 March 7th to 8th detected by the Fermi LAT
17:00 - 17:15	GBM Observations of Terrestrial Gamma-ray Flashes (Foley)
17:15 - 17:30	AGILE observations of Terrestrial Gamma-ray Flashes (Marisaldi)

Esperimenti nello spazio


CONVERSIONE $\gamma \rightarrow e^+ e^-$



- photons materialize into matter-antimatter pairs:
$$E_\gamma \rightarrow m_{e^+}c^2 + m_{e^-}c^2$$
- electron and positron carry information about the direction, energy and polarization of the γ -ray



High-energy γ -ray telescopes work on the principle of pair production. A photon passing through matter may convert into an electron-positron pair.



The probability of such a conversion taking place is roughly independent of the energy of the incident photon above 1 GeV, and falls off at lower energies. While the full pair-production differential cross section is a complex function of incident γ -ray energy, electron and positron energy, nuclear recoil energy, opening angle, azimuthal angle, and recoil angle [1], several simplifying assumptions give simple estimates of bulk behavior [2]. For a homogeneous material the intensity of the incident γ -ray beam falls off like

$$I = I_0 \exp\left(-\frac{7}{9}t/X_0\right), \quad (1)$$


due to all interactions, where t is the thickness of material and X_0 is the radiation length of the material. Therefore, the probability of a particular γ -ray to interact in the material is

$$P(t) = 1 - \exp\left(-\frac{7}{9}t/X_0\right). \quad (2)$$

Pair production offers an opportunity for photons detection. In fact we can estimate the incident γ -ray energy and direction by tracking the resulting e^+e^- pair. The reconstructed energy will be the sum of the e^+ and e^- energies, corrected for energy loss in the instrument, and the incident direction of the γ -ray can be obtained by the momentum-weighted average of the e^+ and e^- directions.

$$\text{angolo di apertura } \theta(e^+ e^-) \approx m_e/k_\gamma$$

DISTRIBUZIONE θ_P




ATIC

4 lanci dal 2000

ATIC must be capable of measuring the incident cosmic ray charge and energy over an energy range of 50 GeV to >100 TeV.


The fully active ATIC calorimeter is composed of 10 layers of Bismuth Germanate (BGO) scintillating crystals and is located on the bottom of the instrument. Above the calorimeter is the target section consisting of three plastic scintillator strip hodoscopes to define the instrument aperture and provide redundant charge and trajectory measurements, as well as layers of inert carbon (between hodoscopes) to provide a volume for the incident particles to interact.

On the top of the detector stack is the highly segmented silicon matrix detector that provides an accurate measure of the incident particle charge. Surrounding the detector stack, electronics bays hold the flight computers, readout electronics, power system boards and other instrument electronics. The total weight of ATIC is about 1,500 kg (3,300 lbs), the total power consumed is less than 350 Watts.





EGRET

COMPTON OBSERVATORY INSTRUMENTS




The Instruments on CGRO Cover Six Orders of Magnitude in Photon Energy



EGRET

- 1991-2000
- 30 MeV - 30 GeV
- AGN, GRB, Unidentified Sources, Diffuse Bkg

AGILE




The AGILE Payload:
the most compact
instrument for high-
energy astrophysics


It combines for the first
time a **gamma-ray
imager** (30 MeV- 30 GeV)
with a **hard X-ray
imager** (18-60 keV) with
large FOVs (1-2.5 sr) and
optimal angular
resolution

LANCIATO NEL 2007 DA UN RAZZO INDIANO DALLA BASE DI SRIHARIKOTA

AGILE



AGILE




AGILE

Table 1 - The AGILE Scientific Performance


Gamma-ray Imaging Detector (GRID)	
Energy range	30 MeV – 50 GeV
Field of view	~ 2.5 sr
Flux sensitivity ($E > 100$ MeV, 5σ in 10^6 s)	3×10^{-7} (ph cm $^{-2}$ s $^{-1}$)
Angular resolution at 100 MeV (68% cont. radius)	3.5 degrees
Angular resolution at 400 MeV (68% cont. radius)	1.2 degrees
Source location accuracy (high Gal. lat., 90% C.L.)	~15 arcmin
Energy resolution (at 400 MeV)	$\Delta E/E \sim 1$
Absolute time resolution	~ 2 μ s
Deadtime	~ 100 – 200 μ s
Hard X-ray Imaging Detector (Super-AGILE)	
Energy range	18 – 60 keV
Single (1-dim.) detector FOV (FW at zero sens.)	$107^\circ \times 68^\circ$
Combined (2-dim.) detector FOV (FW at zero sens.)	$68^\circ \times 68^\circ$
Sensitivity (18-60 keV, 5σ in 1 day)	~ 15 mCrab
Angular resolution (pixel size)	6 arcmin
Source location accuracy (S/N~10)	~1-2 arcmin
Energy resolution (FWHM)	$\Delta E \sim 8$ keV
Absolute time resolution	~ 2 μ s
Mini-Calorimeter	
Energy range	0.35 – 50 MeV
Energy resolution (at 1.3 MeV)	13% FWHM
Absolute time resolution	~ 3 μ s
Deadtime (for each of the 30 CsI bars)	~ 20 μ s

GLAST/FERMI




FERMI Key-Features

- Two GLAST instruments:
 - LAT: 20 MeV - >300 GeV
 - GBM: 10 keV - 25 MeV
 - Launch: 11 June 2008.
 - 565 km, circular orbit
 - 5-year mission (10-year goal)
 - International Collaboration
- Huge field of view:
 - LAT: 20% of the sky at any instant; in sky survey mode, expose all parts of sky for ~30 minutes every 3 hours.
 - GBM: whole unocculted sky at any time.
- Huge energy range, including largely unexplored band 10 GeV- 100 GeV
- *LAT: Large Area Telescope*
- *GBM: Gamma ray Burst Monitor*



FERMI



FERMI Large Area Telescope

- **Precision Si-strip Tracker (TKR)**

18 XY tracking planes. 228 mm pitch). High efficiency. Good position resolution (ang. resolution at high energy) $12 \times 0.03 X_0$ front end => reduce multiple scattering. $4 \times 0.18 X_0$ back-end => increase sensitivity $>1\text{GeV}$. Tot t $\approx 1X_0$

- **CsI Calorimeter(CAL)**

Array of 1536 CsI(Tl) crystals in 8 layers. Hodoscopic => Cosmic ray rejection.

=> shower leakage correction.

$8.5 X_0$ => Shower max contained $<100\text{ GeV}$


- **Anticoincidence Detector (ACD)**

Segmented (89 plastic scintillator tiles)

=> minimize self veto,

Reject background of charged cosmic rays;

- **Electronics System** Includes flexible, robust hardware trigger and software filters.




Height/Width = 0.4

=> Large field of view

Systems work together to identify and measure the flux of cosmic gamma rays with energy 20 MeV - >300 GeV.


GBM



FERMI




FERMI: RISOLUZIONE E




PAMELA

- Costruito in gran parte in Italia
- Lanciato nel 2006 con un razzo russo



PAMELA



PAMELA-MAGNETE




- The magnetic material used is the sintered Nd-Fe-B with a large residual magnetic induction (1.3T). The average field inside the magnet is 0.4 T, with a good homogeneity.
- The combined characteristics of the magnet and of the Si tracker will allow a Maximum Detectable Rigidity (MDR) greater than 740 GV/c.

PAMELA- CALORIMETRO EM




- The total thickness corresponds to 0.9 interaction lengths and 16 radiation lengths.
- The energy resolution for high energy electrons is better than 10% .

AMS




AMS con magnete permanente

$$BL^2=0.14 \text{ Tm}^2$$


AMS-02 Superconducting Magnet Silicon Tracker Layers




AMS-02 Permanent Magnet Silicon Tracker Layers




Layer 9 comes from moving the ladders at the edge of the acceptance from layer 1. The layer 8 is moved on top of the TRD to become 1N.


Risoluzione dello spettrometro



AMS2




Photon conversion



Direct Photon detection

EM sampling calorimeter




- ⇒ **High granularity :**
 - 0.5 Molière radius in X-Y
 - 18 samplings, $0.9 X_0$ in depth


why spaghetti?

- ⇒ best longitudinal & lateral shower reconstruction
 - ⇒ energy correction
 - ⇒ p/e separation

- ⇒ best γ angular resolution



Calorimetro Elettromagnetico




$1X_0$: probabilità $1/e$ di emettere 1γ o e^+e^-

$$\# \text{ fotoni} \propto E \quad \Delta E/E = (a/\sqrt{E}) + b$$


Alcune prestazioni degli esperimenti

SENSIBILITÀ

EGRET(Spark Chamber) VS. GLAST(Silicon Strip Detector)




EGRET on Compton GRO
(1991–2000)




GLAST Large Area Telescope
(2006–2015)

AGILE



Area effettiva di AGILE a 100 MeV confrontata con EGRET in funzione della direzione di incidenza del fotone.

AGILE



Area effettiva in funzione dell'energia del fotone per diversi angoli di incidenza

AMS02 Gamma

Unidentified Sources with AMS


AMS

- Source localization:
 $(E > 10 \text{ GeV}) < 2'$

In 1 Year and for source of strength:
 $5 \times 10^{-8} \text{ ph cm}^{-2} \text{s}^{-1}$ ($E = 1 \text{ GeV}$)

GLAST


- Source localization:
 $< 5'$ and high sensitivity



EGRET


- Source localization:
 $< 30'$
for source of strength
 $10^{-8} \text{ ph cm}^{-2} \text{s}^{-1}$
- Limited sensitivity
above 1 GeV

GLAST: PRESTAZIONI




GLAST

One year All-Sky Survey Simulation, $E\gamma > 100$ MeV




PULSAR: POLAR CAP



Esperimenti Telescopi Cerenkov

SCIAMI IN ATMOSFERA

Cerenkov and Extensive air shower (EAS) gamma ray telescope concepts





CERENKOV IN ATMOSFERA

- A livello del mare $(n - 1) = \varepsilon \approx 3 \cdot 10^{-4}$.
- Per $v \approx c$, $\cos\theta = 1/\beta n \approx 23 \text{ mrad} \approx 1.3^\circ$
- Energia di soglia per l'effetto Cerenkov: $\cos\theta = 1 = 1/\beta n ; \beta > 1/n$
$$E = \gamma mc^2 = mc^2/(1 - \beta^2)^{1/2} ; (1 - \beta^2)^{1/2} = (1 - 1/n^2)^{1/2} = [(n^2 - 1)/n^2]^{1/2}$$
$$E = mc^2/\sqrt{2\varepsilon} \quad 1/\sqrt{2\varepsilon} \approx 41$$
- La soglia per elettroni: $E \approx 21 \text{ MeV}$
muoni: $E \approx 4.4 \text{ GeV}$
- Il massimo di produzione di particelle si ha a 10 km di quota (massimo di produzione Cerenkov).
- L'area illuminata a terra è un ellisse, o un cerchio di raggio $r = h \cdot \theta = 10^4 \cdot 23 \cdot 10^{-3} = 230 \text{ m}$ con una superficie di $1.6 \cdot 10^5 \text{ m}^2$.
- Il numero di fotoni prodotti nel visibile, 350—500 nm, da un gamma di 1 TeV è

$$N\gamma \approx 8.2 \cdot 10^3 \text{ fotoni}/\lambda$$

pari a circa 30—50 fotoni/m² in un'area entro $\approx 100 \text{ m}$ dall'asse dello sciame.



Whipple: diametro=10m, E > 350 GeV

The Whipple collaboration, which pioneered the Imaging Atmospheric Cherenkov Technique for the detection of very high energy (VHE) gamma rays, is based at the Fred Lawrence Whipple Observatory in Southern Arizona, in the United States. **The primary emphasis of the collaboration's research effort is the search for and study of celestial sources of gamma-rays in the energy range of 100 GeV - 10 TeV.**






Fig. 4.— Integral excess events observed by the Whipple telescope from the directions of IH1426+428 (crosses) and the Crab Nebula (open circles) during 2001 as a function of integrated Cherenkov light in the shower image. Exposure on the Crab Nebula was adjusted to match the total excess of IH1426+428 at the lowest size cut applied, 366 digital counts. One photoelectron corresponds to ~ 3.6 digital counts.

MAGIC

230 m²
E: 10-300 GeV
La Palma



MAGIC



MAGIC 24 APRILE 2009



HESS

H.E.S.S. is a next-generation system of Imaging Atmospheric Cherenkov Telescopes for the investigation of cosmic gamma rays in the 100 GeV energy range. The name H.E.S.S. stands for **High Energy Stereoscopic System**, and should also remind of **Victor Hess**, who received in 1936 the **Nobel Prize in Physics for his discovery of cosmic radiation**. The acronym also emphasizes two main features of the proposed installation, namely the **simultaneous observation of air showers with several (3 to 4) telescopes, under different viewing angles**, and the **combination of multiple (up to 16) telescopes to a large system to increase the effective detection area for gamma rays**. With telescopes of over 100 m² mirror area, the proposed system provides a

- detection threshold of about 40 GeV,
- full spectroscopic capability above 100 GeV, an
- angular resolution for individual showers of 0.1 degrees
- energy resolution of about 20%.




It will allow to explore gamma-ray sources with intensities at a level of a few thousandth parts of the flux of the Crab nebula. H.E.S.S. is located in Namibia, near the Gamsberg, an area well known for its excellent optical quality. The first four H.E.S.S. telescopes (Phase I of the H.E.S.S. project) are under construction and are expected to successively go into operation between early 2002 and 2003

I quattro telescopi della Fase I sono stati completati nel dicembre 2003


HESS

$A \approx 100 \text{ m}^2$



HESS

Supernova explosions ...



Le immagini di HESS mostrano chiaramente la shell delle supernove dove hanno origine i gamma



If supernova explosion waves are cosmic accelerators - as long suspected by scientists - they should be clearly visible in gamma rays. Indeed, H.E.S.S. images resolve the ring-like shock fronts for the first time and show them glowing in high-energy gamma rays.

One of the first highlights of the H.E.S.S. observations was the discovery of high-energy gamma rays from two supernova explosion shells, whose 'names' RX J1713.7-3946 and RX J0852.0-4622 refer to their celestial coordinates and the fact that they were first seen as X-ray sources. The H.E.S.S. telescopes were able to resolve the spatial structure of the gamma-ray source; as predicted, gamma rays were found to exactly trace the supernova shell. This discovery proves conclusively that supernova explosion waves work as cosmic accelerators, at least up to energies of 100 million million electrons volts ('electron volt' is a unit characterising the energies of particles and radiation; visible light has 2 to 4 electron volts).

Background

Our entire Galaxy is permeated by cosmic rays - atomic nuclei accelerated to very high energies. The existence of cosmic rays was discovered by Victor Hess in 1912, almost 100 years ago. Throughout this time, the origin of cosmic rays was heavily debated: Somewhere in our Galaxy must be cosmic particle accelerators capable of creating particle energies many orders of magnitude beyond the biggest man-made accelerators on Earth.

HESS: CENTAUROS A



Spectral distribution of the emission from Centaurus A, from X-rays to the VHE gamma-ray energy band. Archival X-ray and EGRET gamma-ray data are shown in grey, previous VHE upper limits and the tentative early detection in purple.



VERITAS




The High Energy Stereoscopic System (H.E.S.S.)



CANGAROO

The “Big Four”




TELESCOPI CERENKOV

Group/ Instrument	Location	Reflector(s) Number × Aperture	Camera Pixels	Threshold (GeV)	Epoch Beginning
<i>Operating Telescopes^a</i>					
Whipple	Arizona, USA	10 m	331	250	1984
Crimea	Crimea, Ukraine	6 × 2.4 m	6 × 37	1000	1985
SHALON	Tien Shen, Russia	4 m	244	1000	1994
CANGAROO	Woomera, Aust.	3.8 m	256	500	1994
HEGRA	La Palma, Sp.	5 × 3 m	5 × 271	500	1994
CAT	Pyrenées	3m	600	250	1996
Durham/ Mark 6	Narrabri, Aust.	3 × 7 m	1 × 109	250	1996
TACTIC	Mt. Abu, India	4 × 3.5 m	1 × 225	300	1997
Seven TA	Utah, USA	7 × 2 m	7 × 256	500	1998
STACEE	Sandia, New Mexico	32 × 7 m	32 × 1	75	1998
CELESTE	Pyrenées, France	40 × 7 m	40 × 1	50	1998
<i>Future Telescopes</i>					
CANGAROO II	Woomera, Aust.	7 m	1 × 512	250	1999
GRAAL/CESA-1	Almeria, Sp.	63 × 7.1 m	4	100	1999
Solar II	Barstow, CA	96 × 7.1 m	96 × 1	20	2002
MAGIC	La Palma, Sp.	17 m	1 × 800	30	2001
HESS	Namibia	4 × 10 m	4 × 700	50	2002
CANGAROO III	Woomera, Aust.	4 × 10 m	4 × 512	75??	2003
VERITAS	Arizona, USA	7 × 10 m	7 × 499	75	2004

^a From Catanese & Weekes 1999

OSSERVAZIONE SPAZIO-TERRA

- Gli esperimenti nello spazio e a Terra sono complementari fra di loro. Nella figura è mostrata la sensibilità al flusso di gamma vs l'energia di GLAST e di HESS. Le linee blu rappresentano rispettivamente il flusso di fotoni dalla Crab Nebula e lo stesso flusso ridotto per un fattore 10 e 100.



RIVELATORI GAMMA

