

Particle Physics - Chapter 12a

LHC – machine and detectors





Paolo Bagnaia
SAPIENZA
UNIVERSITÀ DI ROMA

AA 18-19

12 – LHC – machine and detectors

1. [LHC physics](#)
2. [The LHC Collider](#)
3. [The luminosity](#)
4. [LHC operations](#)
5. [The ATLAS detector](#)
6. [The CMS detector](#)
7. [Detectors comparison](#)
8. [Detector performances](#)
9. [LHC events](#)





LHC physics

- the LHC physics programme still has a long story ahead;
- [the heavy ion programme is outside the scope of the lectures – see ALICE talks]
- until now, its results can be broadly divided into three categories :
 - a. "bread and butter", i.e. quantitative improvements on soft & SM physics;
 - b. **the discovery of the Higgs boson** [*still a tiny probability that the "bump" is NOT the Higgs boson of the SM*];
 - c. searches of physics beyond the SM;
- (a) contains beautiful and intelligent results, from soft physics to jets, from W^\pm/Z to top;
- however, they are too fresh [*imho*] to be part of an institutional course;
- [we all hope that] (c) will be the most interesting;
- however, it is outside the scope of these lectures;
- therefore, this chapter includes two parts :
 1. a general discussion of the method of analysis of LHC, mainly the problems caused by the high \mathcal{L} ;
 2. a report of the Higgs discovery [noblesse oblige];
- the other parts are left to the next semester, your Thesis and (hopefully) your individual research activity.

Enjoy it !



LHC physics: why $10^{34} \text{ cm}^{-2} \text{ s}^{-1}$

Such a large \mathcal{L} is a must or a luxury ?

Compute two toy processes :

- cross section for a s-channel process :

➤ $\sigma \approx K g^2 / s;$

Ex. $\sigma[e^+e^- \rightarrow \gamma^* \rightarrow \mu^+\mu^-] = \frac{4}{3} \pi \alpha^2 / s.$

❖ K : adimensional factor ~ 1
(e.g. $4\pi/3$);

❖ g : coupling constant < 1
(it depends on the dynamics);

❖ s : (energy)² in CM sys;


- formation of a resonance (s-channel)

[e.g. $\sqrt{s} = m_x = 100 \text{ GeV}$]:

❖ $g \sim 10^{-2};$

❖ $m_x \sim 100 \text{ GeV};$



➤ $\sigma \approx K g^2 / m_x^2 =$
 $= [0.389 \text{ GeV}^2 \text{ mb}] \times 10^{-4} / 10^4 \approx$
 $= 4 \times 10^{-36} \text{ cm}^2;$



[of course, it is too simplistic : parton structure functions (pdf), decay BR, detector acceptance, analysis inefficiencies are neglected; but all these effects DECREASE the yield or the identification of the effects.]




LHC physics: plots for $10^{34} \text{ cm}^{-2} \text{ s}^{-1}$



these plots show the trend vs \sqrt{s} of :

- σ_x : s-channel cross section just defined;
 - $\sigma_{\text{tot}}/\sigma_x$: if $\sigma_{\text{tot}} \approx 100 \text{ mb}$, ratio between number of events and interesting ones;
 - lumi@.01Hz : \mathcal{L} to get a rate of .01 Hz for the m_x just defined;
- ∴ obvious, but concerning →

high \mathcal{L} is a must.



LHC physics: events at $10^{34} \text{ cm}^{-2} \text{ s}^{-1}$



How many (interesting) events?
an estimate of the order of
magnitude:

- "average year" $\sim 10^7 \text{ s}$;
- $\mathcal{L}_{\max} \approx 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$;
- $\mathcal{L}_{\text{int}} \approx 10^{41} \text{ cm}^{-2} = 100 \text{ fb}^{-1}$;
- last column roughly includes
the detection efficiencies;
- clearly, it is NOT possible to
record all these events (\rightarrow
act on trigger/selection).

Process	$\sigma (\text{pb})$	rate (@ $10^{34} \text{ cm}^{-2} \text{ s}^{-1}$)	events / year
collisions (bc)	---	4×10^7	4×10^{14}
events	1×10^{11}	1×10^9	10^{16}
$W \rightarrow e\nu$	1.5×10^4	150	10^9
$Z \rightarrow e^+e^-$	1.5×10^3	15	10^8
$t\bar{t}$	800	8	10^8
$b\bar{b}$	5×10^8	5×10^6	10^{13}
$\tilde{g}\tilde{g}$ (SUSY) [$m_g = 1 \text{ TeV}$]	1	0.01	10^5
Higgs [$m_H = 125 \text{ GeV}$]	20	0.2	2×10^6
QCD jets [$p_T > 200 \text{ GeV}$]	10^5	1000	10^{10}

LHC physics: DAQ at $10^{34} \text{ cm}^{-2} \text{ s}^{-1}$

$[\sigma_{\text{tot}}(\text{pp})$ is a fundamental parameter of the Nature; however, here we study it only as an obstacle to observe high- p_T collisions]

- $\mathcal{L} \approx 10^{34} \text{ cm}^{-2} \text{s}^{-1}$ (actually higher);

- $\tau_{\text{bc}} = 25 \text{ ns}$

- $f_{\text{bc}} = 1/\tau_{\text{bc}} = 40 \text{ MHz};$


- $\sigma_{\text{tot}} \approx 100 \text{ mb} (= 10^{-25} \text{ cm}^2);$

- therefore :

- $R_{\text{int}} \approx 1 \text{ GHz};$
- $\mathcal{L}_{\text{bc}} = 2.5 \times 10^{26} \text{ cm}^{-2};$
- $n_{\text{bc}} = 25 \text{ events / bc};$
- $n_{\text{inelast}} \approx 20 \text{ events / bc};$
- $N^{\pm}_{\text{partic.}} \approx 1000 / \text{bc};$
- $dN^{\pm}/d\eta \approx 100 / \text{bc};$
- $W_{\text{detect.}} \approx 3 \text{ kW};$
- $\Delta s_{\text{bc}} = 25 \text{ ns} \times c = 7.5 \text{ m};$



do you see
the paradox ?

- i.e. there are "waves" of $\sim 1000 \pi^{\pm}$ (+ as many γ 's) every 25 ns;
- the waves are on concentric spheres at 7.5 m each other (e.g. at the same time the muon chambers "see" previous bc's respect to the inner detector);
- the detectors must have an adequate bandwidth to cope with it (and the necessary radiation resistance !!!).



LHC physics: trigger at 10^{34} cm $^{-2}$ s $^{-1}$


Trigger Setup



The LHC Collider

150 tonnes Helium, ~90 tonnes at 1.9 K
140 MJ stored beam energy in 2012
450 MJ magnetic energy per sector at 4 TeV


1720 Power converters
> 9000 magnetic elements
7568 Quench detection systems
1088 Beam position monitors
~4000 Beam loss monitors



The LHC Collider: a view



The LHC Collider: the complex



The LHC Collider : parameters


Date	2009	2012	2015	nomin.	Parameter	Value
Maximum beam energy (TeV) ↑	3.5	4	6.5	7	Circumference	26.659 km
Delivered integrated luminosity (fb^{-1}) ↑	up to 5.6	23.3	4	—	Interaction regions	4 total, 2 high \mathcal{L}
Luminosity \mathcal{L} ($10^{33} \text{ cm}^{-2}\text{s}^{-1}$) ↑	3.7	7.7	5.2	>10	Free space at interaction point	38 m
Time between collisions τ_{bc} (ns) ↔	49.90	49.90	24.95	24.95	Magnetic length of dipole	14.3 m
Full crossing angle (μ rad) ↔	240	≈ 300		≈ 300	Length of standard cell	106.9 m
Energy spread $\Delta E/E$ (units 10^{-3}) ↓	0.116	0.116		0.113	Phase advance per cell	90°
Bunch length (cm) ↔	9	9		7.5	Dipoles in ring	1232 main dipoles
Beam radius (10^{-6} m) ↓	26	20		16.6	Quadrupoles in ring	482 2-in-1 + 24 1-in-1
Initial luminosity decay time, $-\mathcal{L}/(d\mathcal{L}/dt)$ (hr) ↑	8	8		14.9	Magnet type	s.c. 2 in 1 cold iron
Transverse emittance ($10^{-9} \pi \text{ rad-m}$) ↓	0.7	0.6		0.5	Peak magnetic field	8.3 T
β^* , ampl. function @ i.p. (m) ↓	1	0.6	0.4	0.55	Injection energy	450 GeV
Beam-beam tune shift / crossing (10^{-4})	23	60		34	RF frequency	400.8 MHz
Particles per bunch (10^{10}) ↑	15	15		11.5		from [PDG]
Bunches per ring per species ↑	1380	1380	2244	2808		
Average beam current / species (mA) ↑	374	374		584		

The LHC Collider: dipoles

1000th Dipole Installed (sep 5, 2007)




The LHC Collider: dipole structure



The LHC Collider: dipole operations

Mike Lamont – CERN academic training – sep 2013
(read it ! it's wonderful !!!)



Dipoles

- Number 1232
- Field (450 GeV) 0.535 T
- Field (7 TeV) 8.33 T
- Bending radius 2803.95 m
- Main Length 14.3 m


Horizontal force component per quadrant (nominal field)
1.7 MN/m.

Force tends to "open" the magnet, hence the Austenitic steel collars.


[more info : <http://lhc-machine-outreach.web.cern.ch/lhc-machine-outreach/>]

The LHC Collider: injection cycle

from Mike Lamont




The LHC Collider: data-taking cycle



The LHC Collider: nominal cycle

Globally the machine state is fairly well described by machine mode/beam mode combination



Mike Lamont

The LHC Collider: the user view

LHC Page1

Fill: 2514

E: 4000 GeV

t(SB): 00:57:54

14-04-12 17:13:53

PROTON PHYSICS: STABLE BEAMS

Energy:

4000 GeV

I(B1):

1.31e+14

I(B2):

1.32e+14

intensity 1

intensity 2


energy

luminosity 1

luminosity 2

luminosity ...

status



Comments 14-04-2012 16:19:52 :

BIS status and SMP flags

B1 B2

Link Status of Beam Permits

Global Beam Permit

Setup Beam

Beam Presence

Moveable Devices Allowed In

Stable Beams

true

true

false

false

true

true

true

true

true

true

true

true

*** STABLE BEAMS ***

AFS: 50ns_1092b_1051_0_1032_108bpi12inj

PM Status B1

ENABLED


PM Status B2

ENABLED




The luminosity: \mathcal{L}_{int} vs time

CMS Integrated Luminosity Delivered, pp





The luminosity: \mathcal{L}_{int} vs time

LHC performances 2017




An almost impossible achievement: efficiency > 90% for many months [the hole is just a period of LHC maintenance].




"what really counts is not the immediate act of courage or of valor, but those who bear the struggle day in and day out - not the sunshine patriots but those who are willing to stand for a long period of time", John F. Kennedy.

The luminosity: $\mathcal{L}_{\text{peak}}$

- In 2016 LHC has achieved the luminosity foreseen in the project, i.e. $\mathcal{L} = 10^{34} \text{ cm}^{-2}\text{s}^{-1}$...
- ... and in 2017-18 it doubled it ($\mathcal{L} = 2 \times 10^{34} \text{ cm}^{-2}\text{s}^{-1}$);
- for $\sqrt{s} = 14 \text{ TeV}$, wait another couple of years.
- [1 Hz/nb = $10^{33} \text{ cm}^{-2}\text{s}^{-1}$]



The luminosity : $\langle n_{\text{int}} \rangle$



$$\mu = \langle n_{\text{int}} \rangle = \mathcal{L} \tau_{\text{bc}} \sigma_{\text{inel}} = [\text{e.g. } \approx]$$

$$\approx (10^{34}) \times (25 \times 10^{-9}) \times (8 \times 10^{-26}) \approx 20.$$

see § 8

Pros and cons of the value of μ at LHC:

☺ for fixed τ_{bc} , $\mu \propto \mathcal{L}$, so large μ necessary for rare processes, like Higgs;

- ☺ for fixed \mathcal{L} , $\mu \propto \tau_{\text{bc}}$; so a decrease in μ is payed by a decrease in τ_{bc} , the processing time for the trigger and DAQ (now 25 ns, the bare minimum);
- ☹ large $\mu \rightarrow$ many overlapping events
→ systematics in trigger thresholds;
→ systematics in vertex reconstruction;
→ systematics in calo calibrations and reconstruction;
→ mistakes in assignment of heavy flavors, jets, muons to event;
→ (... many other problems ...)
- ☺ some of the LHC data have been taken with a different τ_{bc} (50 ns instead of 25 ns); for the same \mathcal{L} , this fact doubles μ (\rightarrow 25 ns is better than 50 ns, but ...)
- ☺ anyway, **large μ is necessary**, so you better learn to survive with it.



LHC operations: 2018...

LHC schedule 2018

(the last year of operations
before LS2, see later)

$> 65 \text{ fb}^{-1}$



keeping the LHC availability
close to 50% (stable beams)

CERN

www.cern.ch
 Status of LHC machine and plans
 Frédéric Bordry
 ATLAS week
 19th February 2018

LHC 2018 Operations Summary Schedule													
Wk	Jan 1	Jan 2	Jan 3	Jan 4	Jan 5	Jan 6	Jan 7	Jan 8	Jan 9	Jan 10	Jan 11	Jan 12	Jan 13
Mo	1	8	15	22	29	5	12	19	26	5	12	19	26
Tu													
We													
Th													
Fr													
Sa													
Su													
Start powering tests													
LHC, T12, T18 closed													
Machine checkout													
Start Beam Commissioning													
Wk	14	15	16	17	18	19	20	21	22	23	24	25	26
Mo	Easter	2	9	16	Scrubbing	23	30	7	14	Whit Sun	21	28	4
Tu						1st May							
We												TS1	
Th					Interleaved commissioning & intensity ramp up			Ascension					8 th = 90 m run
Fr											MD 1		VdM program
Sa													
Su													
First Stable beams													
Collisions with 1200 bunches													
Wk	27	28	29	30	31	32	33	34	35	36	37	38	39
Mo	8 th = 90 m run	2	9	16	23	30	6	13	20	27	3	10	17
Tu													
We					MD 2								TS2
Th											Jeune G.		
Fr											MD 3		
Sa													
Su													
End of run [xsec]													
Wk	40	41	42	43	44	45	46	47	48	49	50	51	52
Mo	1	8	15	22	MD 4	28	5	12	19	26	3	10	Xmas 24
Tu													
We					Special physics run			MD 5					
Th						TS3							
Fr													
Sa													
Su					MD 4								
Powering Tests / Magnet Training													
Long Shutdown 2													

LHC operations: 2015-2023




LHC operations: HL-LHC




- March 2016: HL-LHC included in the ESFRI (European Strategy Forum on Research Infrastructures) roadmap as "landmark project" in March 2016.
- June 2016: HL-LHC project formally approved by CERN's Council.
- "Full exploitation of the LHC physics potential with the HL-LHC phase is the top priority of the ESPP [European Strategy for Particle Physics] and the highest near-term large-project priority of the US P5 roadmap." "LHC/HL-LHC is CERN's flagship project for the next 20 years." [Fabiola Gianotti, CERN's Scientific Strategy, ECFA HL-LHC Experiments Workshop, Aix-Les-Bains, 3/10/2016].


LHC operations: HL-LHC performances




The ATLAS detector



The ATLAS detector: scheme



The ATLAS detector: inner tracker




Pixel	SCT	TRT
3 cylindrical layers	4 cylindrical layers	73 straw planes
2x3 disks	2x9 disks	160 straw planes



The ATLAS detector: calorimeters

sandwich
scint-Fe


"accordion"
LAr - Pb



The ATLAS detector: e.m. calo




e/ γ




- "accordion" LAr – Pb
- cryogenic
- hermetic
- longitudinal + radial segmentation

The ATLAS detector: e^\pm id and measure




an electron is detected many (> 100) times after the interaction point; even the non-detection in the had. calo is important (cfr a γ in the pixels/SCT/TRT).



The ATLAS detector: had. calo



- "tiles" Fe – Scintillator
- WLS readout
- hermetic
- high segmentation



The ATLAS detector: μ spectrometer








The ATLAS detector: μ chambers




Schematic view of:

- A. a chamber of drift tubes
- B. a single tube
- C. a muon hitting a tube
- D. the hit time distribution
- E. the r-t relation



The CMS detector



The CMS detector: view

CMS DETECTOR

Total weight : 14,000 tonnes

Overall diameter : 15.0 m

Overall length : 28.7 m

Magnetic field : 3.8 T

STEEL RETURN YOKE
12,500 tonnes

SILICON TRACKERS

Pixel ($100 \times 150 \mu\text{m}$) $\sim 16\text{m}^2$ $\sim 66\text{M}$ channels

Microstrips ($80 \times 180 \mu\text{m}$) $\sim 200\text{m}^2$ $\sim 9.6\text{M}$ channels

SUPERCONDUCTING SOLENOID
Niobium titanium coil carrying $\sim 18,000\text{A}$

MUON CHAMBERS

Barrel: 250 Drift Tube, 480 Resistive Plate Chambers


Endcaps: 468 Cathode Strip, 432 Resistive Plate Chambers

PRESHOWER
Silicon strips $\sim 16\text{m}^2$ $\sim 137,000$ channels


FORWARD CALORIMETER
Steel + Quartz fibers $\sim 2,000$ Channels

CRYSTAL
ELECTROMAGNETIC
CALORIMETER (ECAL)
 $\sim 76,000$ scintillating PbWO_4 crystals




HADRON CALORIMETER (HCAL)
Brass + Plastic scintillator $\sim 7,000$ channels



The CMS detector: scheme






The CMS detector: inner tracker



Si pixel + strip
detector



The CMS detector: e.m. calo



e.m. calo:
 PbWO_4 crystals

The CMS detector: had. calo


HCAL (tower structure):

- Barrel (HB): $|\eta| < 1.4$, 2304 towers
- End caps (HE): $1.3 < |\eta| < 3.0$, „ towers
- Outside coil (HO): $|\eta| < 1.26$ (tail catcher)
→ 4608 towers (Plastic scintillator tiles, $\approx 10 \lambda_N$)
→ $\Delta\eta \times \Delta\phi \approx 0.087 \times 0.087 \rightarrow 0.350 \times 0.175$

- Forward (HF): $2.9 < |\eta| < 5.0$ (not shown)
- 2 x 900 towers (Quartz fibers, $\approx 10 \lambda_N$)
- $\Delta\eta \times \Delta\phi \approx 0.111 \times 0.175 \rightarrow 0.302 \times 0.350$


CASTOR calorimeter (not shown):

- $5.1 < |\eta| < 6.5$, $\approx 22 X_0$, $\approx 10 \lambda_N$



hadronic calo:
Brass/scintillator/wls readout


The CMS detector: μ system



muon system:

drift tube (DT) chambers

Detectors comparison : ATLAS vs CMS



Detectors comparison : structure

*thanks to
Anna Colaleo*

	 ATLAS	 CMS
Magnet(s)	Air-core toroids + Solenoid in inner cavity Calorimeters outside field 4 magnets	Solenoid Calorimeters inside field 1 magnet
Tracker/ Inner Detector	Silicon pixels, Silicon strips, Transition Radiation Tracker. 2T magnetic field	Silicon pixels, Silicon strips. 4 T magnetic field
Electro-magnetic calorimeter	Lead plates as absorbers with liquid argon as the active medium	Lead tungstate (PbW04) crystals both absorb and respond by scintillation
Hadronic calorimeter	Iron absorber with plastic scintillating tiles as detectors in central region, copper and tungsten absorber with liquid argon in forward regions.	Stainless steel and copper absorber with plastic scintillating tiles as detectors
Muon detector	Large air-core toroid magnets with muon chamber form outer part of the whole ATLAS	Muons measured already in the central field, further muon chambers inserted in the magnet return yoke

Detectors comparison : performances



thanks to
Anna Colaleo

	 ATLAS	 CMS
Tracker/ Inner Detector	TRD → particle identification $\sigma/p_T \approx 5 \times 10^{-4} p_T$ (GeV) $\oplus 0.01$	No particle identification $\sigma/p_T \approx 1.5 \times 10^{-4} p_T$ (GeV) $\oplus 0.005$
Electro-magnetic calorimeter	$\sigma/E \approx 10\%/\sqrt{E}$ (GeV) Longitudinal segmentation	$\sigma/E \approx (2 \div 5)\%/\sqrt{E}$ (GeV) No longitudinal segmentation
Hadronic calorimeter	$> 10 \lambda$ $\sigma/E \approx 50\%/\sqrt{E}$ (GeV) $\oplus 0.03$	$> 5.8 \lambda + \text{tail catcher}$ $\sigma/E \approx 65\%/\sqrt{E}$ (GeV) $\oplus 0.05$
Muon detector	air $\sigma/p_T \approx 7\% @ 1 \text{ TeV}$ (spectrometer alone)	Fe $\sigma/p_T \approx 5\% @ 1 \text{ TeV}$ (combining spectrometer + tracker)

imho (*common, but not unanimous*):

- two complementary strategies almost everywhere;
- ... with different optimizations (e.g. resolution vs robustness);
- a textbook example of "guided" detector design;
- ... to guarantee optimal results (\rightarrow not miss major discoveries).

Detectors comparison : mag. spectrometers



ATLAS:


- main magnet: solenoid $B = 0.7 \text{ T}$;
- bending in (r,z) ;
- straight tracks in (r,ϕ) ;
- at small r , a solenoid $B = 2 \text{ T} \rightarrow$ bending also in (r,ϕ) ;
- less precise in extrapolating to main vtx;
- μ -system in air \rightarrow no multiple scatt. for μ 's;
- larger bending for μ at large $\eta \rightarrow$ more precise.

CMS:


- main magnet: solenoid $B = 4 \text{ T}$;
- bending in (r,ϕ) ;
- straight tracks in (r,z) ;
- more precise in extrapolating to main vtx;
- μ -system in Fe \rightarrow large multiple scatt. for μ 's;
- less bending for μ 's at large η .

*thanks to
Anna Colaleo*


Detector performances : $Z \rightarrow e^+e^-$




- the classic benchmark of tracker + e.m. calo.;
- no way to improve wrt LEP, used only for detector debug/calibration.



Detector performances : $Z \rightarrow \mu^+\mu^-$




- the classic benchmark of tracker + muon chambers;
- [no way to improve wrt LEP, used only for detector debug/calibration.]




[time evolution of physics analyses :
discovery \rightarrow precision meas \rightarrow
detector study \rightarrow background]




Detector performances : $J/\psi \rightarrow \mu^+\mu^-$



- $Z \rightarrow \mu^+\mu^-$ and $J/\psi Z \rightarrow \mu^+\mu^-$ are ideal channels for μ studies :
- inner detector + muon spectrometer;
 - agreement ($\text{MC} \leftrightarrow \text{data}$) \rightarrow confidence in analysis (including errors !).





Detector performances : silicon trackers




- resolution of few μm necessary for impact parameter \rightarrow identification of secondary vertexes \rightarrow heavy flavors \rightarrow higgs;
- agreement (MC \leftrightarrow data) \rightarrow confidence in analysis (including errors !).

Detector performances : vertex resolution

- Vertex resolution as a function of the its track multiplicity.
- Expect $\sigma \approx \sigma_0/\sqrt{N} + \sigma_1$;
- σ_0 and σ_1 constants, different for xy (transverse) and z (beam axis).




Detector performances : e.m. calo



- although real life is somewhat different (sys from cell-to-cell calib, control of temperature, etc), test-beam results are impressive;
- expect $\sigma/E \approx \sigma_1/E \oplus \sigma_2 / \sqrt{E} \oplus \sigma_3$;
- σ_1 looks negligible, while σ_3 dominates at high E.

→ test these expectations with real particles:



- Z (previous slides);
- π^0, η, \dots (next slides)



Detector performances : $\pi^0, \eta \rightarrow \gamma\gamma$

The π^0 and η widths are a measurement of the electro calo resolution in a difficult environment (inside jets or in high multiplicity events).


Notice the good (almost perfect) agreement with MC predictions.






Detector performances : jet response

jet resolution as a function of p_T^{jet} :

- measured for different event types;
- stat and (mainly) syst uncertainty
2%, almost independent on p_T .




Detector performances : E_T



$E_T [= |\sum \vec{E}_j^T|]$ for events with e^\pm or μ^\pm with $p_T > 20 \text{ GeV}$:

- great agreement with predictions;
- reliable measurement

Detector performances: ATLAS μ^\pm



$\Delta p_T / p_T$ vs p_T [project, low η] :

▼ meas. error + calib ($\propto p_T$);

○ chamber alignment ($\propto p_T$);

□ multiple scattering ($\propto \approx \text{const}$);


○ $\Delta E_\mu(\text{calo})$ fluctuations (tail at high loss measurable from brem shower);

○ at spectrometer entrance
(= ▼ \oplus ○ \oplus □);


△ total at main vertex (= ○ \oplus ○).

- at low p_T ($p_T < 200$ GeV) vtx extrapolation (○) and scattering (□) give the main contributions;
- at high p_T the accuracy of the spectrometer (▼ \oplus ○) dominates;
- at fixed p_T and high η (not shown), Δp_T gets worse.

Detector performances : mass($\mu^+\mu^-$)



Detector performances: $W^\pm \rightarrow e^\pm\nu$




Detector performances: trigger thresholds


- e^+e^- : small cross section $\rightarrow [R = \mathcal{L}\sigma \approx \text{few Hz}] \rightarrow \text{event trigger}$, i.e. trigger on single bunch crossing, if it contains an event candidate; @ LEP, $1-\varepsilon \approx 10^{-3}$, negligible dead time;
- $p\bar{p}(\bar{p}\bar{p})$: high hadronic total cross section $\rightarrow [R = \mathcal{L}\sigma \approx 10^6 - 10^9 \text{ Hz}] \rightarrow$ rates too big (and uninteresting events) $\rightarrow \text{physics trigger}$, i.e. select a (tiny) fraction of events, which exhibit peculiar

characteristics (i.e. high- p_T , multileptons, high E_T ...); use cuts (i.e. thresholds), user defined in kinematical variables;

- the thresholds are applied on a kinematical variable "x" (e.g. p_T^{lepton}), measured in a rough and fast way by the trigger detector(s); therefore the experimenters have to compromise among rejection, efficiency, dead time, bandwidth ... and physics.




Detector performances: μ -trigger lvl-1



Efficiency ϵ at level 1 :

- vs p_T : notice the "size" of the threshold;
- vs η , integrated for $p_T > 14$ GeV : notice the flatness.




NB the effective yield $N_{\text{obs}} = N_{\text{produced}} * \epsilon$:

- the bulk of the data is near p_T threshold
- ... where ϵ is varying;
- ... and the physics less interesting.


Detector performances: μ -trigger HLT

Efficiency ε vs p_T at the highest trigger level (HLT):


- notice the sharper "size" of the threshold (\rightarrow less useless data);
- ... at the price of a much higher threshold (\rightarrow no recovery of events lost in lvl1);
- ... with the advantage of (much) smaller rates :
 $O(10 \text{ KHz}) @ \text{lvl-1} \rightarrow O(10 \text{ Hz})$.



LHC events : Pb Pb \rightarrow Z X \rightarrow e⁺e⁻ X



LHC events : 78 primary interactions





LHC events : 2 jets, $p_T \approx 2$ TeV




Run Number: 201006, Event Number: 55422459


Date: 2012-04-09 14:07:47 UTC



LHC events : a multijet event




LHC events : $W \rightarrow e\nu$



CMS Experiment at LHC, CERN
Run 133874, Event 21466935
Lumi section: 301
Sat Apr 24 2010, 05:19:21 CEST


Electron $p_T = 35.6 \text{ GeV}/c$
 $M_{ET} = 36.9 \text{ GeV}$
 $M_T = 71.1 \text{ GeV}/c^2$




LHC events : $H \rightarrow ZZ^* \rightarrow (e^+e^-)(\mu^+\mu^-)^*$

2e2 μ candidate with $m_{2e2\mu} = 123.9$ GeV

$p_T(e, e, \mu, \mu) = 18.7, 76, 19.6, 7.9$ GeV, $m(e^+e^-) = 87.9$ GeV, $m(\mu^+\mu^-) = 19.6$ GeV



LHC events : $H \rightarrow ZZ^* \rightarrow (\mu^+\mu^-)(\mu^+\mu^-)^*$




Run Number: 189280,
Event Number: 143576946
Date: 2011-09-14, 11:37:11 CE'

EtCut>0.3 GeV
PtCut>3.0 GeV
Vertex Cuts:
Z direction <1cm
Rphi <1cm

Muon: blue
Cells: Tiles, EMC

LHC events : $H \rightarrow W^+W^- \rightarrow e^+\nu\mu^-\nu$




LHC events : $Z \rightarrow \mu^+\mu^-$, $Z \rightarrow \mu^+\mu^-$


Run: 338220

Event: 2718372349



2017-10-15 00:50:49 CEST



LHC events : $H \rightarrow \gamma\gamma$



LHC events : $H \rightarrow \gamma\gamma$



(c) CERN. All rights reserved.

References: collider & experiments

1. LHC : JINST 3 (2008) S08001.
2. LHC : L.Evans, Ann. Rev. Nucl. Part. Sci. 2011. 61:435–66.
3. LHC (recent) : J. Wenninger, PoS (Charged 2018) 001.
4. ATLAS detector : JINST 3 (2008) S08003.
5. ATLAS events :
<https://twiki.cern.ch/twiki/bin/view/AtlasPublic/EventDisplayPublicResults/>
6. CMS detector : JINST 3 (2008) S08004.
7. CMS events : <https://cdsweb.cern.ch/>
8. [see also references on results]



Jacopo Robusti (Tintoretto) - The Forge of Vulcan, ca. 1578
Palazzo Ducale - Venice



SAPIENZA
UNIVERSITÀ DI ROMA

End of chapter 12