


## Resonances and synchronization

by Guido Gentile, Alessandro Giuliani, GG, [arXiv:1106.1476](#)

- 1) Quasi integrable systems
  - 2) Chaotic systems
- 

$$H(\vec{A}, \vec{a}) = \frac{1}{2}\vec{A}^2 + \varepsilon f(\vec{A}, \vec{a})$$



Representation of phase space in terms of  $\ell$  rotators:

$$\vec{a} = (a_1, \dots, a_\ell) \in T^\ell, \quad \vec{A} = (A_1, \dots, A_\ell)$$

Unperturbed system  $\Rightarrow$  motions will have **all** spectra

$$\bar{A} = \bar{A}_0, \quad \bar{a} = \bar{a}_0 + \bar{\omega}t, \quad \bar{\omega} = (\omega_1, \dots, \omega_\ell)$$

$\Rightarrow$  in particular  $\omega_j$  **rationally depend.**: e.g.  $(\omega_1^0, \dots, \omega_{\ell'}^0, 0, 0, \dots)$

Such motions are called **resonant**: more generally

$$\begin{cases} \bar{A} = \bar{A}_0 + \bar{X}(\bar{\psi}) \\ \bar{a} = R\bar{\psi} + \bar{Y}(\bar{\psi}) \end{cases}, \quad \bar{\psi} \in T^{\ell'}, \quad \ell' < \ell$$

$\bar{X}, \bar{Y}$  smooth  $R \ell \times \ell'$  integer matrix and the

$$\bar{\psi} \rightarrow \bar{\psi} + \bar{\omega}^0 t$$


give a solution to the eq. of motion then: **resonant**.

If  $\ell' = 1$  periodic: since Poincaré;  $\ell' > 1$

If  $1 < \ell' < \ell$  KAM-theory: mild conditions **but non trivial**

Example  $\vec{a} = (\vec{a}_1, \vec{a}_2) \in T^{\ell'} \times T^{\ell-\ell'}$ ,  $f(\vec{a}) \equiv f(\vec{a}_1, \vec{a}_2)$

$X_\varepsilon(\vec{\psi}), Y_\varepsilon(\vec{\psi})$  analytic in  $\varepsilon$ ,  $\vec{\psi}$  exist with domain including



$$\bar{f}(\vec{a}_2) = \int \frac{d\vec{a}_1}{(2\pi)^{\ell'}} f(\vec{a}_1, \vec{a}_2)$$

$$\partial \bar{f}(\vec{a}_2^0) = \vec{0}, \quad \partial^2 \bar{f}(\vec{a}_2^0) < 0$$

$$|\vec{\omega}^0 \cdot \vec{v}| > C|\vec{v}|^{-\tau}, \text{ Diophantine p.}$$

Resonances exist **at real  $\varepsilon$**  points

(Llave-Zhou, Gentile-G)

“Intrinsic res”. “Extrinsic” res. or **synchronization**:


$$\begin{cases} \dot{\vec{a}} = \vec{A} \\ \dot{\vec{A}} = -\varepsilon \partial_{\vec{a}} V(\vec{a}) + \varepsilon \vec{F}(\vec{\omega}t) \end{cases}$$

$\exists$  motion with spectrum  $\vec{\omega}$  ?  $\ell = 2$  (Corsi-Gentile). Friction?

# Chaotic systems: paradigm Anosov flow periodically forced

- 1) volume preserving
- 2) dissipative


**No quasi periodic** motions and **few periodic** ones: finitely many with period less than any  $T < \infty$ . So in this case only extrinsic resonances properly can exist: **synchronization**.




$$Sx = \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$$

$$\dot{x} = \delta(z)(Sx - x)$$

$$\dot{w} = 1 \quad \dot{z} = 1$$






$$\dot{x} = \delta(z)(Sx - x) + \varepsilon f_\varepsilon(x, w, z)$$

$$\dot{w} = 1 + \varepsilon g_\varepsilon(x, w, z)$$

$$\dot{z} = 1$$




Look at Poincaré's map  $\mathcal{S}$  at  $t = 2\pi n$

$$x' = Sx + \varepsilon \bar{f}_\varepsilon(x, w)$$

$$w' = w + \varepsilon \bar{g}_\varepsilon(x, w)$$

a) volume preserving (if  $\varepsilon = 0$  the  $\mathcal{S}$  is not even ergodic & has one “central” Lyapunov exponent 0)

**Th.** *as close as wished (in  $C^2$ ) to  $\bar{f} = \bar{g} = 0 \exists$  open set of perturbations with  $\mathcal{S}_\varepsilon$*



(1) *Ergodic*

(2) *Central Lyap. exp.  $\ell_\varepsilon > 0$*

(3)  *$\exists \mathcal{S}$ -invariant foliation  $\Lambda$  into  $C^1$ -smooth lines  $l$*

(4)  *$\exists k$  and  $E$  of full vol. s.t.  $E \cap l$  is exactly  $k < \infty$  pts (!)*

[Conjectures:  $k > 1$  &  $k = 1$ ]

*SW 1999, RW 2001*

No synchronization: in  $(x, w, z)$  the planes  $w = \text{const}$  are invariant under the P.-map but **volatilize** under perturbation

Is synch. possible in dissipation? which attractor structure?

1) Simplest possibility attractor=“periodic orbit” on a orbit close to an unperturbed periodic one.

2) An attractor of “pathological nature” like the volume preserving cases but with Hausdorff dimension lower.

3) **A periodic strange attractor: dissipation stabilizes a single one among the unperturbed invariant surfaces  $w = const$**

This can be easily tested in simulations: a variety of phenomena show up: consistent with 2) or 3). “Naivest” case

$$f = 0, \quad g(x, w, z) = (\sin(z - w) + \sin(x_1 + z + w))$$


(first studied) immediately shows an instance of (3). Special:

$$g_0(x, w) \stackrel{\text{def}}{=} \int_0^{2\pi} g(x, w + t, t) dt,$$
$$g_1(x, w) \stackrel{\text{def}}{=} \int_0^{2\pi} \frac{\partial}{\partial w} g(x, w + t, t) dt$$

Notice that in example  $g_0(x, \pi) = 0$ ,  $g_1(x, \pi) = \Gamma < 0$ ,  $\forall x$ .

**Th.** *If  $\exists w_0, 0 < \varepsilon_0$  such that*

*$g_0(x, w_0) = 0$ ,  $g_1(x, w_0) = \Gamma < 0$ ,  $\forall x$  then for  $0 < \varepsilon < \varepsilon_0$*



*(1)  $\exists$  attractor  $w(x) = w_0 + U_\varepsilon(x)$*

*(2)  $U(x)$  is  $h$ -Hölder continuous*

$$h \geq \frac{|\Gamma|}{\log \beta_+} \varepsilon + O(\varepsilon^2)$$

*(3)  $U_\varepsilon$  is analytic in  $\varepsilon$  in  $\mathcal{D}$*

Assumptions can be relaxed into

(a)  $\int_0^{2\pi} dt g(x, w_0 + t, t) = \varepsilon \bar{g}(x)$ , for some  $\bar{g}(x)$ ,

(b)  $\int_0^{2\pi} dt \partial_w g(x, w_0 + t, t) \leq \Gamma$ , for  $\Gamma < 0$ ,

**Conjectures**

(a')  $\int_0^{2\pi} dt g(x, w_0 + t, t) = \tilde{g}(x)$ , with  $\tilde{g}(x)$  with 0 average,

(b')  $\int_0^{2\pi} dt \partial_w g(x, w_0 + t, t) = \tilde{g}_1(x)$ , with  $\tilde{g}_1(x)$  with  $< 0$  average.



## Basics

$$(x(t), w(t), t) = (x, w + t + u(x, t), t), \quad t \in (0, 2\pi]$$

$$\begin{aligned} \dot{x} &= \delta(z)(Sx - x) + \varepsilon f_\varepsilon(x, w, z) & u(x, 0) &= U(x) \\ \dot{w} &= 1 + \varepsilon g_\varepsilon(x, w, z) & u(x, 2\pi) &= U(Sx) \\ \dot{z} &= 1 \end{aligned}$$

Taylor expansion in  $u$  to second order

$$\begin{aligned} \dot{u}(x, t) &= \varepsilon g(x, w + t + u(x, t), t) \\ &\equiv \varepsilon g(x, w_0 + t, t) + \varepsilon \partial_w g(x, w_0 + t, t) u(x, t) + \varepsilon G(x, t, u(x, t)) \end{aligned}$$

Solved “as a linear equation” in terms of “**Wronskian**”

$$\begin{aligned} \Gamma(x, t, \tau) &\stackrel{\text{def}}{=} \int_\tau^t \partial_w g(x, w + y, y) dy: \\ u(x, t) &= e^{\varepsilon \Gamma(x, t, 0)} u(x, 0) \\ &+ \int_0^t e^{\varepsilon \Gamma(x, t, \tau)} (\varepsilon g(x, w_0 + \tau, \tau) + \varepsilon G(x, \tau, u(x, \tau))) d\tau \end{aligned}$$

*Ideas: equations and invariance condition* ( $\Gamma(x, 2\pi, 0) \equiv \Gamma < 0$ )

$$\begin{aligned}u(x, t) &= e^{\varepsilon\Gamma(x,t,0)} u(x, 0) \\ &+ \int_0^t e^{\varepsilon\Gamma(x,t,\tau)} (\varepsilon g(x, w_0 + \tau, \tau) + \varepsilon G(x, \tau, u(x, \tau))) \\ U(Sx) &= e^{\varepsilon\Gamma} U(x) \\ &+ \int_0^{2\pi} e^{\varepsilon\Gamma(x,2\pi,\tau)} (\varepsilon g(x, w_0 + \tau, \tau) + \varepsilon G(x, \tau, u(x, \tau))) d\tau\end{aligned}$$

The assumptions  $\Gamma < 0$  and  $\int_0^{2\pi} g(x, w_0 + t, t) dt = 0$  imply

$$|u(\cdot, 2\pi)|_\infty < e^{\varepsilon\Gamma} |u(\cdot, 0)|_\infty + O(\varepsilon^2) + O(\varepsilon|u|_\infty^2) \leq e^{\frac{1}{2}\varepsilon\Gamma} |u|_\infty$$

if  $\frac{\delta}{2} < |u(x, 0)| < \delta$  with  $\delta$  small and  $0 \leq \varepsilon \ll \delta$ .

Hence **there is an attractor in the slab  $[w_0 - \delta, w_0 + \delta]$ : but why is it a surface?**

Next idea: **replace** some  $\varepsilon$ 's with  $\mu$ :

$$u(x, t) = e^{\mu\Gamma(x,t,0)} u(x, 0) + \int_0^t e^{\mu\Gamma(x,t,\tau)} (\varepsilon g(x, w_0 + \tau, \tau) + \varepsilon G(x, \tau, u(x, \tau))) d\tau$$
$$U(Sx) = e^{\mu\Gamma} u(x) + \int_0^{2\pi} e^{\mu\Gamma(x,2\pi,\tau)} (\varepsilon g(x, w_0 + \tau, \tau) + \varepsilon G(x, \tau, u(x, \tau))) d\tau$$

- 1) **Fix**  $\mu$  small and **prove** existence of  $U(x)$  analytic in  $|\varepsilon| < C(\mu)$  complex.
- 2) Study  $C(\mu)$  and **show**  $C(\mu) > c\sqrt{\mu}$ .
- 3) Conclude by  $\varepsilon = \mu$

Why Hölder continuity? convergence in  $\varepsilon$  reduces the question to first order. It is explicitly evaluated and shows the property of **tiny Hölder continuity**,  $O(\varepsilon)$ .